

THE STRONG SYMMETRIC GENUS OF THE HYPEROCTAHEDRAL GROUPS

MICHAEL A. JACKSON

ABSTRACT. The strong symmetric genus $\sigma^0(G)$ is the smallest genus of a surface on which G acts faithfully as a group of orientation preserving automorphisms. In this paper we give the strong symmetric genus for the hyperoctahedral groups.

1. INTRODUCTION

In the study of orientable surfaces, it is natural to consider the finite group, which is the group of automorphisms of a given orientable surface. On the other hand, given a finite group, one may consider the surfaces on which the group acts faithfully. There are several natural parameters assigned to a group, which are associated to the action of that group on compact surfaces. The first of these parameters is the genus $\gamma(G)$ of the group G , which is the smallest genus of a surface on which some Cayley graph for G can be embedded. The symmetric genus $\sigma(G)$ of a finite group G is the smallest genus of a surface on which G acts faithfully as a group of automorphisms. The strong symmetric genus $\sigma^0(G)$ is the smallest genus of a surface on which G acts faithfully as a group of orientation preserving automorphisms. (See [10] chapter 6.) It is clear that these parameters satisfy the inequality $\gamma(G) \leq \sigma(G) \leq \sigma^0(G)$.

Perhaps the most classical of these parameters is the strong symmetric genus. It was first considered by Burnside [2], who included the further restriction that the quotient space under the action be a sphere. The strong symmetric genus is what we will concern ourselves with in this paper. Various results have been shown involving the strong symmetric genus. For example, all groups G such that $\sigma^0(G) \leq 3$ have been computed (see [1, 11]). Also the strong symmetric genus of several infinite families of finite groups have been found, such as the alternating and symmetric groups [3, 4, 5] and the groups $PSL_2(p)$ [9]. In addition, the strong symmetric genus has been found for the sporadic finite simple groups [6, 17, 18, 19].

Since the strong symmetric genus of the alternating and symmetric groups is known, it is natural to try to find the strong symmetric genus for other infinite families of finite groups. An example of such an infinite family is the collection of hyperoctahedral groups. Recall that the hyperoctahedral groups are the finite Coxeter groups of type B_n . The hyperoctahedral group B_n for $n \geq 3$ is defined as the group of symmetries of the n -dimensional cube. In this paper we will prove the following theorem about the hyperoctahedral groups:

Theorem 1. *For all $n \geq 3$, except $n = 5, 6$, and 8 , the strong symmetric genus of the hyperoctahedral group B_n is $\frac{n!2^n}{24} + 1 = \frac{n!2^{n-3}}{3} + 1$. The strong symmetric genus in the exceptional cases B_5, B_6 , and B_8 are 289, 3841, and 645,121 respectively.*

The investigation of the strong symmetric genus of the hyperoctahedral groups grew out of a VIGRE Working Group at The Ohio State University during the fall of 2002. In this working group, we were looking to find the strong symmetric genus of some finite Coxeter groups. The exceptional cases of Theorem 1 were found by the working group.

2. SUBGROUPS OF THE HYPEROCTAHEDRAL GROUPS

For the present time we will fix an $n \geq 3$ and look at the hyperoctahedral group B_n . As we have previously said, B_n is defined as the group of symmetries of the n -dimensional cube. This group is also described as the wreath product $\mathbb{Z}_2 \wr \Sigma_n$. Another description of this group is as the group of all $n \times n$ signed permutation matrices. For our purposes, we will use the wreath product description as adopted by V. S. Sikora [12]. For an element of B_n , we will write a tuple $[\sigma, b]$ where σ is an element of Σ_n and b is a list of n binary digits representing the element of $(\mathbb{Z}_2)^n$. The multiplication then becomes $[\sigma, b] \cdot [\tau, c] = [\sigma \cdot \tau, \tau^{-1}(b) + c]$ where addition in the binary digits is a parity computation. We will use the convention of calling b even or odd according to the number of ones appearing as binary digits of b . Notice that if b and c have the same parity, then $b + c$ is even, and if they differ in parity, then $b + c$ is odd.

Recall that the following sequence is a split exact sequence of groups:

$$(\mathbb{Z}_2)^n \xrightarrow{i} B_n \xrightarrow{\pi} \Sigma_n$$

where $i(b) = [1, b]$ and $\pi([\sigma, b]) = \sigma$. We will also assume that $s : \Sigma_n \rightarrow B_n$ with $\pi \circ s = id_{\Sigma_n}$. Proper subgroups $G \subset B_n$ such that $\pi(G) = \Sigma_n$ are of particular interest here. Specifically we wish to examine maximal subgroups of B_n with this property. Throughout this discussion, we will assume that $n \geq 5$. In order to examine the maximal subgroups of B_n with $\pi(G) = \Sigma_n$, we need to introduce another maximal subgroup of B_n . Let C_n be the subgroup of B_n consisting of all elements $[\sigma, b]$ such that $\sigma \in A_n$. It is immediately clear that $[B_n : C_n] = 2$ and $C_n = \mathbb{Z}_2 \wr A_n$. So we get the following commutative diagram

$$\begin{array}{ccccc} (\mathbb{Z}_2)^n & \xrightarrow{i} & C_n & \xrightarrow{\pi} & A_n \\ \parallel & & \downarrow & & \downarrow \\ (\mathbb{Z}_2)^n & \xrightarrow{i} & B_n & \xrightarrow{\pi} & \Sigma_n \end{array}$$

where both horizontal sequences are split exact.

Let $G \subset B_n$ be a subgroup such that $\pi(G) = \Sigma_n$. Let $H = G \cap C_n$. We, thus, see that $\pi(H) = A_n$ and $[G : H] = 2$. Letting $K = \ker(\pi|_H)$ gives the following

commutative diagram:

$$\begin{array}{ccccc}
K & \longrightarrow & H & \xrightarrow{\pi} & A_n \\
\downarrow & & \downarrow & & \parallel \\
(\mathbb{Z}_2)^n & \xrightarrow{i} & C_n & \xrightarrow{\pi} & A_n
\end{array}$$

Notice that C_n acts on $(\mathbb{Z}_2)^n$ by conjugation with kernel $i((\mathbb{Z}_2)^n)$. Since $K \triangleleft H$, we see that K must be invariant under the action of A_n . The image of K , therefore, must be one of the following cases:

- (1) $K \cong (\mathbb{Z}_2)^n$,
- (2) $K = 1$,
- (3) $K \cong \mathbb{Z}_2$, or
- (4) $K \cong (\mathbb{Z}_2)^{n-1}$.

We will look at each of the above cases. The first case obviously implies that $H = C_n$ and $G = B_n$. The second case clearly implies $H \cong A_n$ and $G \cong \Sigma_n$. In the third case, we see that $K = Z(C_n)$ and so $H \cong \mathbb{Z}_2 \times A_n$ and $G \cong \mathbb{Z}_2 \times \Sigma_n$. The fourth case is the most interesting one for this discussion. Notice that $[C_n : H] = 2$, so $H \triangleleft C_n$. Let s be any splitting homomorphism of the split exact sequence defining C_n . Then $s(A_n) \cap H \triangleleft s(A_n)$ and $[s(A_n) : H \cap s(A_n)] \leq 2$. Since $n \geq 5$, A_n is simple, so $s(A_n) \subset H$. We have just proven the following proposition:

Proposition 2. *For $n \geq 5$, let H be a subgroup of C_n with $\pi(H) = A_n$; then H is a split extension of A_n by one of the following: 1 , \mathbb{Z}_2 , $(\mathbb{Z}_2)^{n-1}$, or $(\mathbb{Z}_2)^n$. In addition in the first two cases, H is isomorphic to A_n and $\mathbb{Z}_2 \times A_n$ respectively. In the last case, $H = C_n$, and in the third case $H = \{[\sigma, b] \in C_n \mid b \text{ is even}\}$.*

We now use the following commutative diagram to relate Proposition 2 to the group G :

$$\begin{array}{ccccc}
K & \longrightarrow & H & \xrightarrow{\pi} & A_n \\
\parallel & & \downarrow & & \downarrow \\
K & \longrightarrow & G & \xrightarrow{\pi} & \Sigma_n \\
\downarrow & & \downarrow & & \parallel \\
(\mathbb{Z}_2)^n & \xrightarrow{i} & B_n & \xrightarrow{\pi} & \Sigma_n
\end{array}$$

Straightforward calculations using the fact that $[G : H] = 2$ gives us the following proposition:

Proposition 3. *For $n \geq 5$, let G be a subgroup of B_n with $\pi(G) = \Sigma_n$; then G is a split extension of Σ_n by one of the following: 1 , \mathbb{Z}_2 , $(\mathbb{Z}_2)^{n-1}$, or $(\mathbb{Z}_2)^n$. In addition in the first two cases, G is isomorphic to Σ_n and $\mathbb{Z}_2 \times \Sigma_n$ respectively. In the last case*

$G = B_n$, and in the third case either

$$G = \{[\sigma, b] \in B_n | b \text{ is even}\} \text{ or } G = \left\{ [\sigma, b] \in B_n \left| \begin{array}{l} b \text{ is even if } \sigma \in A_n \\ b \text{ is odd if } \sigma \in \Sigma_n \setminus A_n \end{array} \right. \right\}.$$

3. GENERATORS IN B_n

In order to find the strong symmetric genus of a group, it will be important to find a pair of generators. Of particular interest will be the orders of the generators in this pair as well as the order of their product. In the following proposition, we will show that in the case of the hyperoctahedral groups, these orders will all be even.

Proposition 4. *If $[\sigma, b]$ and $[\tau, c]$ generate B_n , then the orders of $[\sigma, b]$, $[\tau, c]$, and their product $[\sigma \cdot \tau, \tau^{-1}(b) + c]$ all are even.*

Proof: Suppose that one of the orders is odd. Without loss of generality, we may assume that $[\sigma, b]$ has odd order. Clearly σ and τ must generate Σ_n . Since $[\sigma, b]$ has odd order, σ must also have odd order; therefore, $\sigma \in A_n$, $\tau \in \Sigma_n \setminus A_n$, and $\sigma \cdot \tau \in \Sigma_n \setminus A_n$. So we see that τ and $\sigma \cdot \tau$ have even orders.

Let k be the order of σ . Since $[\sigma, b]$ has odd order, $[\sigma, b]^k = 1$. Notice that b must be even since $[\sigma, b]^k = [1, \sigma^{-k+1}(b) + \sigma^{-k+2}(b) + \dots + \sigma^{-1}(b) + b]$ and an addition of an odd number of odd elements is odd. If c is also even, then $[\sigma, b]$ and $[\tau, c]$ both lie in the proper subgroup $\{[\sigma, b] \in B_n | b \text{ is even}\} \subset B_n$ and, therefore, cannot generate B_n . If on the other hand c is odd, then $[\sigma, b]$ and $[\tau, c]$ both lie in the proper subgroup

$$\left\{ [\sigma, b] \in B_n \left| \begin{array}{l} b \text{ is even if } \sigma \in A_n \\ b \text{ is odd if } \sigma \in \Sigma_n \setminus A_n \end{array} \right. \right\} \subset B_n$$

and, therefore, cannot generate B_n . Thus the order of $[\sigma, b]$ must also be even. \square

Notice that if $[\sigma, b]$ and $[\tau, c]$ generate B_n , then σ and τ must generate Σ_n . Since we are looking to find generators of B_n , we need to first find generators of Σ_n . The second step in the process is to construct generators of B_n from the generators of Σ_n . Also during this construction we would like to have control of the orders of the generators of B_n as well as their product. The following proposition gives the results of the process we will use to construct generators of B_n from those of Σ_n .

Proposition 5. *Suppose σ and τ generate Σ_n as $\text{Sym}(\Gamma)$ where $\Gamma = \{1, 2, \dots, n\}$ such that $\sigma \cdot \tau \in \Sigma_n \setminus A_n$ where σ , τ , and $\sigma \cdot \tau$ all have even order. Furthermore, assume that σ fixes an element $i \in \Gamma$ and τ fixes at least three elements of Γ , one of which we call j such that i and j are in the same cycle of the element $\sigma \cdot \tau$. Let $b = (0, \dots, 0, 1, 0, \dots, 0)$ where the 1 is in the i^{th} position, and let $c = (0, \dots, 0, 1, 0, \dots, 0)$ where the 1 is in the j^{th} position. Under these conditions, $[\sigma, b]$ and $[\tau, c]$ generate B_n . In addition the elements $[\sigma, b]$, $[\tau, c]$ and $[\sigma \cdot \tau, \tau^{-1}(b) + c]$ have the same orders as σ , τ , and $\sigma \cdot \tau$ respectively.*

Proof: The results about the orders of $[\sigma, b]$ and $[\tau, c]$ are obvious since $\sigma(b) = b$ and $\tau(c) = c$. On the other hand, $(\sigma \cdot \tau)^{-k}(\tau^{-1}(b) + c) = (\sigma \cdot \tau)^{-k-1}(b) + (\sigma \cdot \tau)^{-k}(c)$.

Notice that if k is the length of the cycle in $\sigma \cdot \tau$, which contains i and j , then $[\sigma \cdot \tau, \tau^{-1}(b) + c]^k = [(\sigma \cdot \tau)^k, (0, 0, \dots, 0, 0)]$ since the images of b will put a 1 in each position corresponding to the elements in the cycle and the images of c will put a 1 in each position corresponding to the elements in the same cycle. Thus it is clear that the order of $[\sigma \cdot \tau, \tau^{-1}(b) + c]$ is the same as the order of $\sigma \cdot \tau$.

Let $G = \langle [\sigma, b], [\tau, c] \rangle \subset B_n$. We need to show that $G = B_n$. Notice that G is a subgroup of B_n such that $\pi(G) = \Sigma_n$. So from Proposition 3 we know that G is a split extension of Σ_n by one of the following: 1, \mathbb{Z}_2 , $(\mathbb{Z}_2)^{n-1}$, or $(\mathbb{Z}_2)^n$.

Recall that any section homomorphism $s : \Sigma_n \rightarrow B_n$ takes $\alpha \in A_n$ to $[1, d] \cdot [\alpha, (0, \dots, 0)] \cdot [1, d]^{-1}$ for some $[1, d] \in B_n$, and if $\alpha \in \Sigma_n \setminus A_n$, then either $s(\alpha) = [1, d] \cdot [\alpha, (0, \dots, 0)] \cdot [1, d]^{-1}$ or $s(\alpha) = [1, d] \cdot [\alpha, (1, \dots, 1)] \cdot [1, d]^{-1}$. Notice if $\alpha \in A_n$, $s(\alpha) = [\alpha, a]$ where a is even.

Suppose first that G is a split extension of Σ_n by 1. Then $[\sigma, b]$ and $[\tau, c]$ must be in the image of some section homomorphism $s : \Sigma_n \rightarrow B_n$. Either $\sigma \in A_n$ or $\tau \in A_n$, however, and both b and c are odd. So G is not a split extension of Σ_n by 1.

Suppose that G is a split extension of Σ_n by \mathbb{Z}_2 . This implies that for some section homomorphism, $s : \Sigma_n \rightarrow B_n$, $G = Z(B_n) \times s(\Sigma_n)$. If n is even, then the only elements in $Z(B_n) \times s(\Sigma_n)$ of the form $[\alpha, a]$ with $\alpha \in A_n$ have an even a . Clearly either $[\sigma, b] \notin Z(B_n) \times s(\Sigma_n)$ or $[\tau, c] \notin Z(B_n) \times s(\Sigma_n)$; therefore, we may assume that n is odd. Suppose that $\tau \in A_n$. So $[\tau, c] = [1, (1, \dots, 1)] \cdot [1, d] \cdot [\tau, (0, \dots, 0)] \cdot [1, d]^{-1}$ for some $[1, d] \in B_n$. It follows that $c = (1, \dots, 1) + \tau^{-1}(d) + d$, which cannot be the case since τ fixes an element besides j . So we may also assume that $\tau \in \Sigma_n \setminus A_n$.

Since $\tau \in \Sigma_n \setminus A_n$, either $[\tau, c] = [1, (1, \dots, 1)] \cdot [1, d] \cdot [\tau, (0, \dots, 0)] \cdot [1, d]^{-1}$ or $[\tau, c] = [1, d] \cdot [\tau, (1, \dots, 1)] \cdot [1, d]^{-1}$ for some $[1, d] \in B_n$. In either case, $c = (1, \dots, 1) + \tau^{-1}(d) + d$, which as before cannot be the case since τ fixes an element besides j . So G is not a split extension of Σ_n by \mathbb{Z}_2 .

Suppose that G is a split extension of Σ_n by $(\mathbb{Z}_2)^{n-1}$. By Proposition 3, either

$$G = \{[\sigma, b] \in B_n \mid b \text{ is even}\} \text{ or } G = \left\{ [\sigma, b] \in B_n \mid \begin{array}{l} b \text{ is even if } \sigma \in A_n \\ b \text{ is odd if } \sigma \in \Sigma_n \setminus A_n \end{array} \right\}.$$

Recall that either $\sigma \in A_n$ or $\tau \in A_n$. If $\sigma \in A_n$, then $[\sigma, b] \notin G$. If $\tau \in A_n$, then $[\tau, c] \notin G$. So G is not a split extension of Σ_n by $(\mathbb{Z}_2)^{n-1}$.

The only possibility left is that G is a split extension of Σ_n by $(\mathbb{Z}_2)^n$. Thus we see that $G = B_n$ and so the proposition is proven. \square

4. GENERATING PAIRS

If a finite group G has generators x and y of orders p and q respectively with xy having the order r , then we say that (x, y) is a (p, q, r) generating pair of G . By the obvious symmetries concerning generators, we will use the convention that $p \leq q \leq r$. Following the convention of Marston Conder in [5], we say that a (p, q, r) generating

pair of G is a minimal generating pair if there does not exist a (k, l, m) generating pair for G with $\frac{1}{k} + \frac{1}{l} + \frac{1}{m} > \frac{1}{p} + \frac{1}{q} + \frac{1}{r}$.

Recall that the groups of small strong symmetric genus are well known (see [1, 11]). So for now we will assume that $\sigma^0(G) > 1$. The hyperoctahedral groups have strong symmetric genus at least 2. It is also known that for groups with $\sigma^0(G) > 1$, any generating pair will be a (p, q, r) generating pair with $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} < 1$. From any generating pair of G , we get an upper bound on the strong symmetric genus of G . If G has a (p, q, r) generating pair, then

$$\sigma^0(G) \leq 1 + \frac{1}{2}|G| \cdot \left(1 - \frac{1}{p} - \frac{1}{q} - \frac{1}{r}\right).$$

The following lemma, which is a result of Singerman [13] (see also [11, 14]), shows that the strong symmetric genus for most groups is computed directly from a minimal generating pair.

Lemma 6 (Singerman [13]). *Let G be a finite group such that $\sigma^0(G) \geq 2$. If $|G| > 12(\sigma^0(G) - 1)$, then G has a (p, q, r) generating pair with*

$$\sigma^0(G) = 1 + \frac{1}{2}|G| \cdot \left(1 - \frac{1}{p} - \frac{1}{q} - \frac{1}{r}\right).$$

In addition, we may assume that either $p = 2$, $(p, q, r) = (3, 3, 4)$, or $(p, q, r) = (3, 3, 5)$.

Applying Lemma 6 to our situation with the hyperoctahedral groups results in the following corollary.

Corollary 7. *Let $n \geq 3$ and $G = B_n$. If $|G| > 12(\sigma^0(G) - 1)$, then G has a (p, q, r) generating pair with*

$$\sigma^0(G) = 1 + \frac{1}{2}|G| \cdot \left(1 - \frac{1}{p} - \frac{1}{q} - \frac{1}{r}\right).$$

In addition, we may assume that $p = 2$, $q = 4$, and r is either 6, 8 or 10.

Proof: Since all groups H with $\sigma^0(H) < 2$ have been classified, we know that $\sigma^0(G) \geq 2$. We also know that if G has a (p, q, r) generating pair, then p , q , and r must all be even. The rest of the corollary follows from straightforward calculations. \square

We will see that the exceptional cases of Theorem 1 will be shown by using GAP [8]. In light of Corollary 7, proving the main theorem for the remaining cases has been reduced to showing that each hyperoctahedral group has a $(2, 4, 6)$ generating pair. We will show the existence of a $(2, 4, 6)$ generating pair for B_n by constructing it from a $(2, 4, 6)$ generating pair for Σ_n using Proposition 5.

5. SPECIAL GENERATORS OF Σ_n

First we will need to recall a theorem of Jordan (see [15]).

Theorem 8 (Jordan [15]). *Let G be a transitive permutation group on a set of size n such that G is primitive and contains a p -cycle for some prime $p < n - 2$. Then G is either A_n or Σ_n .*

We will use this theorem by applying the following corollary. The method that we will use follows that of Marston Conder (see [3, 4]).

Corollary 9. *Let $G = \langle \sigma, \tau \rangle$ be a transitive permutation group on a set Γ of size n such that G contains a p -cycle for some prime $p < n - 2$, which contains an element $a \in \Gamma$ as well as its image under σ and an element $b \in \Gamma$ as well as its image under τ . Then G is either A_n or Σ_n .*

Proof: Using Theorem 8, the only part we need to prove is that G is a primitive permutation group on Γ . Suppose that G acts imprimitively on Γ . Let z be the p -cycle in the hypothesis. Since z fixes all the other points of Γ , the p points permuted by z must lie in the same block of imprimitivity B . By hypothesis, however, a and $\sigma(a)$ are both in B ; therefore, σ fixes B . Also by hypothesis, b and $\tau(b)$ are both in B ; therefore, τ fixes B . This implies that the block B is fixed by all of G , contradicting the transitivity of G . \square

At this point we have done all of the preliminary work; it is now time to prove the main theorem. To begin with we point out that the result for the groups B_n for $3 \leq n \leq 40$ have been shown using GAP [8]. For the groups B_n for $n \geq 41$, we show that each group Σ_n for $n \geq 41$ can be generated by two elements σ and τ . Using Corollary 9, this process will involve finding two elements σ and τ such that these two elements generate a transitive permutation group on n letters such that some element of $\langle \sigma, \tau \rangle$ is a p -cycle, which contains a letter and its image under σ and a possibly different letter and its image under τ . In order to be able to construct a (2,4,6) generating pair, $([\sigma, b], [\tau, c])$, of B_n from σ and τ as in Proposition 5, we will require that the following hold:

- (1) σ has order 4 and τ has order 2,
- (2) σ fixes a letter i ,
- (3) τ fixes at least three letters one of which we call j ,
- (4) $\sigma\sigma \cdot \tau \in \Sigma_n \setminus A_n$,
- (5) i and j are in the same cycle of the element $\sigma \cdot \tau$,
- (6) $\langle \sigma, \tau \rangle$ is a transitive permutation group on n letters, and
- (7) some element of $\langle \sigma, \tau \rangle$ is a p -cycle, which contains a letter and its image under σ and a possibly different letter and its image under τ .




FIGURE 1. Example Σ_{19}




FIGURE 2. Σ_n for $n \equiv 41(12)$ and $n \geq 41$

6. COSET DIAGRAMS AND RESULTS

We will demonstrate the existence of the desired elements σ and τ in each Σ_n for $n \geq 41$ by using coset diagrams. First let us give a brief introduction to coset diagrams by looking at an example in Figure 1. The diagram in Figure 1 is a representation of the group Σ_{19} . The diagram gives two elements of a transitive permutation group on 19 letters. The first element σ is given by the squares and circles in the diagram and will have order 4. A cycle of σ is given by reading the labels in a counter-clockwise direction. So in this case,

$$\sigma = (1, 2, 3, 4)(5, 6, 7, 8)(9, 10, 11, 12)(13, 14)(15, 16, 17, 18).$$

The second element τ is given by the other lines and curves in the diagram. τ will be an involution with a 2-cycle, for each line or curve, that is the cycle of its two labels.




FIGURE 3. Σ_n for $n \equiv 42(12)$ and $n \geq 42$




FIGURE 4. Σ_n for $n \equiv 43(12)$ and $n \geq 43$

In this case we see

$$\tau = (1, 15)(2, 5)(3, 4)(7, 17)(8, 9)(12, 13)(14, 16)(18, 19).$$

To find to where an integer is mapped under the element $\sigma \cdot \tau$, we start at that label, trace along a square or circle in the counter-clockwise direction if possible, and then trace a curve not in one of these shapes if possible. The permutation given by a general word in σ and τ can be found in a similar way.




FIGURE 5. Σ_n for $n \equiv 44(12)$ and $n \geq 44$




FIGURE 6. Σ_n for $n \equiv 45(12)$ and $n \geq 45$

We see that the element $\sigma^2 \cdot \tau$ has four 4-cycles and one 3-cycle, which is $(10, 13, 12)$. By Corollary 9, we see that σ and τ generate the symmetric group on 19 elements. Using the elements σ and τ and creating elements $[\sigma, b]$ and $[\tau, c]$ of B_{19} as in Proposition 5, we also see that $[\sigma, b]$ and $[\tau, c]$ generate B_{19} .

In a similar way as the example above, we will now look at Figures 2 through 13. These figures give coset diagrams for the groups Σ_n for $41 \leq n \leq 52$. As with the




FIGURE 7. Σ_n for $n \equiv 46(12)$ and $n \geq 46$




FIGURE 8. Σ_n for $n \equiv 47(12)$ and $n \geq 47$

example in Figure 1, these figures show that the groups B_n for $41 \leq n \leq 52$ are all $(2, 4, 6)$ groups. The necessary information about $\langle \sigma, \tau \rangle$ containing a p -cycle, which contains a letter and its image under σ and a possibly different letter and its image under τ , can be found in the table in Figure 15.

To show the result for the groups B_n for $n > 52$ are all $(2, 4, 6)$ groups, we will recursively add the piece shown in Figure 14. We attach the piece by matching up the points a and b and then making the a' and b' into a new a and b respectively.




FIGURE 9. Σ_n for $n \equiv 48(12)$ and $n \geq 48$




FIGURE 10. Σ_n for $n \equiv 49(12)$ and $n \geq 49$

Attaching this piece recursively to each of the twelve original diagrams gives a diagram for each $n \geq 41$. Again the necessary information about $\langle \sigma, \tau \rangle$ containing a p -cycle, which contains a letter and its image under σ and a possibly different letter and its image under τ , can be found in the table in Figure 15. Each $n \geq 41$ satisfies the conditions of at least one of the rows of the table. The word in σ and τ in that row then has some power which is a p -cycle for the prime p in the last column. The infinite collection of diagrams created by our recursion together with the table in Figure 15




FIGURE 11. Σ_n for $n \equiv 50(12)$ and $n \geq 50$




FIGURE 12. Σ_n for $n \equiv 51(12)$ and $n \geq 51$

show that every B_n for $n \geq 41$ is a $(2, 4, 6)$ group. This finishes the proof of the main theorem. \square




FIGURE 13. Σ_n for $n \equiv 52(12)$ and $n \geq 52$




FIGURE 14. additional piece

REFERENCES

- [1] S. Allen Broughton, Classifying finite group actions on surfaces of low genus, *J. Pure Appl. Algebra* 69 (1991) 233-270.
- [2] W. Burnside, *Theory of groups of finite order*, Cambridge University Press, 1911.
- [3] Marston D. E. Conder, Generators for alternating and symmetric groups, *J. London Math. Soc.* (2) 22 (1980) 75-86.
- [4] Marston D.E. Conder, More on generators for alternating and symmetric groups, *Quart. J. Math. Oxford* (2) 32 (1981) 137-163.
- [5] Marston D.E. Conder, Some results on quotients of triangle groups, *Bull. Austral. Math. Soc.* 29 (1984) 73-90.
- [6] M.D.E. Conder, R.A. Wilson, A.J. Woldar, The symmetric genus of sporadic groups, *Proc. Amer. Math. Soc.* 116 (1992) 653-663.
- [7] Brent Everitt, Permutation representations of the $(2, 4, r)$ Groups, *Bull. Austral. Math. Soc.* 49 (1994) 499-511.

n value	conditions on m		word in σ and τ	prime
$n = 41 + 12m$	$m \not\equiv 0(13)$		$\sigma\tau\sigma\tau\sigma$	13
$n = 41 + 12m$	$m \not\equiv 8(13)$	$m \not\equiv 9(13)$	$\sigma^3\tau\sigma\tau\sigma\tau\sigma^2\tau$	13
$n = 42 + 12m$	$m \not\equiv 1(7)$		$\sigma\tau\sigma^3\tau$	7
$n = 42 + 12m$	$m \not\equiv 2(7)$	$m \not\equiv 4(7)$	$\sigma^3\tau\sigma\tau\sigma^2\tau\sigma\tau\sigma^2\tau\sigma^3\tau\sigma^2$	7
$n = 43 + 12m$	$m \not\equiv 4(7)$		$\tau\sigma\tau\sigma^3$	7
$n = 43 + 12m$	$m \not\equiv 2(7)$	$m \not\equiv 3(7)$	$\tau\sigma^2\tau\sigma\tau\sigma\tau\sigma^2\tau\sigma\tau\sigma^2\tau\sigma\tau$	7
$n = 44 + 12m$	$m \not\equiv 1(13)$		$\tau\sigma\tau\sigma^3\tau\sigma\tau\sigma^3\tau\sigma$	13
$n = 44 + 12m$	$m \not\equiv 9(13)$	$m \geq 1$	$\sigma^2\tau\sigma^2\tau\sigma\tau\sigma^3\tau\sigma\tau\sigma\tau\sigma^3$	13
$n = 45 + 12m$	$m \not\equiv 5(7)$	$m \not\equiv 6(7)$	$\sigma^2\tau\sigma^2\tau\sigma$	7
$n = 45 + 12m$	$m \not\equiv 1(7)$	$m \not\equiv 3(7)$	$\tau\sigma^3\tau\sigma\tau\sigma\tau\sigma^3\tau\sigma^2$	7
$n = 46 + 12m$	$m \not\equiv 0(7)$		$\tau\sigma^3\tau\sigma^2\tau\sigma\tau\sigma^2\tau\sigma^3\tau\sigma^2\tau\sigma\tau\sigma^3\tau\sigma\tau$	7
$n = 46 + 12m$	$m \not\equiv 1(7)$	$m \not\equiv 5(7)$	$\tau\sigma^2\tau\sigma^3\tau\sigma\tau\sigma\tau\sigma^3\tau\sigma$	7
$n = 47 + 12m$	$m \not\equiv 5(7)$	$m \not\equiv 6(7)$	$\sigma^2\tau\sigma\tau\sigma\tau\sigma^3\tau$	7
$n = 47 + 12m$	$m \not\equiv 1(7)$	$m \not\equiv 3(7)$	$\sigma\tau\sigma^2\tau\sigma\tau\sigma\tau\sigma^2\tau\sigma^3\tau$	7
$n = 48 + 12m$	$m \not\equiv 1(13)$		$\sigma\tau\sigma\tau\sigma$	13
$n = 48 + 12m$	$m \not\equiv 11(13)$	$m \geq 1$	$\sigma\tau\sigma\tau\sigma^2\tau\sigma^3\tau$	13
$n = 49 + 12m$	$m \not\equiv 5(7)$	$m \not\equiv 6(7)$	$\sigma^2\tau\sigma\tau\sigma\tau\sigma^3\tau$	7
$n = 49 + 12m$	$m \not\equiv 1(7)$	$m \not\equiv 3(7)$	$\sigma^2\tau\sigma\tau\sigma^3\tau\sigma\tau\sigma^3\tau\sigma\tau\sigma^2\tau\sigma$	7
$n = 50 + 12m$	$m \not\equiv 4(7)$		$\tau\sigma\tau\sigma\tau\sigma^2\tau\sigma^2\tau\sigma\tau\sigma\tau$	7
$n = 50 + 12m$	$m \not\equiv 6(7)$	$m \geq 2$	$\sigma^2\tau\sigma\tau\sigma^3\tau\sigma\tau\sigma^2\tau\sigma^3\tau\sigma^3$	7
$n = 51 + 12m$	$m \not\equiv 6(7)$		$\sigma^3\tau\sigma^2\tau\sigma\tau\sigma\tau\sigma^2\tau\sigma^2\tau$	7
$n = 51 + 12m$	$m \not\equiv 3(7)$		$\sigma\tau\sigma^2\tau\sigma\tau\sigma^2\tau\sigma\tau\sigma^3\tau\sigma\tau\sigma^3$	7
$n = 52 + 12m$	$m \not\equiv 6(11)$		$\tau\sigma^2\tau\sigma\tau\sigma^2\tau\sigma^3$	11
$n = 52 + 12m$	$m \not\equiv 10(11)$	$m \geq 1$	$\sigma^2\tau\sigma^2\tau\sigma\tau\sigma\tau\sigma^2$	11

FIGURE 15. Demonstration of p -cycles

- [8] The GAP Group, GAP — Groups, Algorithms, and Programming, Version 4.2; Aachen, St Andrews, 1999. (<http://www-gap.dcs.st-and.ac.uk/~gap>)
- [9] Henry H. Glover, Denis Sjerve, Representing $PSL_2(p)$ on a Reiman surface of least genus, Enseignement Math. 31 (1985) 305-325.
- [10] Jonathan L. Gross, Thomas W. Tucker, Topological graph theory, Wiley-Interscience 1987.
- [11] Coy L. May, Jay Zimmerman, Groups of small strong symmetric genus, J. Group Theory 3 (2000) 233-245.
- [12] V.S. Sikora, Two-element bases of hyperoctahedral groups, Visn. Kiiiv. Univ. Ser. Fiz.-Mat. Nauki 1 (1999) 87-93.
- [13] David Singerman, Symmetries of Riemann surfaces with large automorphism group, Math. Ann. 210 (1974) 17-32.
- [14] Thomas W. Tucker, Finite groups acting on surfaces and the genus of a group, J. Combin. Theory Ser. B 34 (1983) 82-98.
- [15] H. Wielandt, Finite permutation groups, Academic Press, London and New York, 1964.

- [16] Arthur White, *Graphs, groups and surfaces*, North-Holland Mathematical Studies 8, North-Holland, 1973.
- [17] Robert A. Wilson, The Monster is a Hurwitz group, *J. Group Theory* 4 (2001) 376-374.
- [18] Robert A. Wilson, The symmetric genus of the Fischer group Fi_{23} , *Topology* 36 (1997) 379-380.
- [19] Robert A. Wilson, The symmetric genus of the Baby Monster, *Quart. J. Math. Oxford Ser. (2)* 44 (1993) 513-516.